

LIGHT VECTOR MESON PHOTOPRODUCTION AT HIGH $|t|$

D. YU. IVANOV

*Institute of Mathematics,
630090 Novosibirsk, Russia
E-mail: d-ivanov@math.nsc.ru*

We have studied in perturbative QCD all independent helicity amplitudes describing the photoproduction of light vector mesons at large t . We found a new hard production mechanism which is related to the possibility for a real photon to fluctuate into a massless $q\bar{q}$ pair in a chiral-odd spin configuration. Each helicity amplitude is given as a sum of chiral-even contribution (when the helicities of quark and antiquark are antiparallel) and this additional chiral-odd part (where the helicities of quark and antiquark are parallel). The chiral-odd contribution is large, it leads to a dominance of the non spin-flip amplitude in a very broad region of intermediately high $|t|$.

1 Introduction

My talk is based on our recent publication¹. We derived pQCD predictions for helicity amplitudes of the high $|t| = -\mathbf{q}^2$ diffractive light vector meson photoproduction process. Initial proton disintegrates into hadron system X which is separated from the produced vector meson V by the rapidity gap η_0 . The cross section for this reaction can be related to those for the photoproduction of V off a quark and a gluon via the gluon and quark densities in a proton $G(x, t)$ and $q(x, t)$:

$$\frac{d^2\sigma(\gamma p \rightarrow VX)}{dt dx} = \sum_f (q(x, t) + \bar{q}(x, t)) \frac{d\sigma(\gamma q \rightarrow Vq)}{dt} + G(x, t) \frac{d\sigma(\gamma G \rightarrow VG)}{dt}; \quad x = \frac{4|t|}{s} \cosh^2 \frac{\eta_0}{2}. \quad (1)$$

There are three independent helicity amplitudes for the parton subprocess $\gamma q \rightarrow Vq$, the first and the second indices are the helicities of photon and vector meson respectively

$$M_{++}(M_{--} = M_{++}), \quad M_{+0}(M_{-0} = -M_{+0}), \quad M_{+-}(M_{-+} = M_{+-}).$$

Within usual perturbation theory a photon can split only into a chiral-even (the helicities of the quark and the antiquark are antiparallel) massless quark pair. The violation of chiral symmetry, which is well known to be a soft QCD phenomenon, generates a nonperturbative chiral-odd component of the real

photon wave function. The interaction of this additional chiral contribution can however be described in pQCD since high t quark-dipole scattering chooses a $q\bar{q}$ configuration with small transverse interquark distances. As a result the chiral-odd contribution can be factorized into a convolution of two nonperturbative photon and vector meson light-cone wave functions with the hard scattering amplitude. The chiral-odd wave function of a real photon have a similar form as the chiral-odd wave function of a vector meson. The photon dimensional coupling constant f_γ (wich is similar to vector meson chiral-odd constant f_V^T) is a product of the quark condensate $\langle \bar{q}q \rangle$ and its magnetic susceptibility²,

$$f_\gamma = \langle \bar{q}q \rangle \chi \approx 70 MeV. \quad (2)$$

Both parameters $\langle \bar{q}q \rangle$ and χ describing QCD vacuum have been tested in various QCD sum rule applications.

2 Results and Discussion

The helicity amplitudes of the process $\gamma q \rightarrow Vq$ are the sums of the chiral-even and the chiral-odd contributions

$$M_{\lambda_1 \lambda_2} = M_{\lambda_1 \lambda_2}^{even} + M_{\lambda_1 \lambda_2}^{odd}.$$

At asymptotically high $|t|$ the dominant helicity amplitude is M_{+0} . Its chiral-even part M_{+0}^{even} has the minimal, $\sim 1/q^3$, suppression.

$$M_{+0}^{even} = -is\alpha_s^2 \frac{32\pi}{3\sqrt{2}} eQ_V f_V/q^3; \quad (3)$$

We present our results for helicity amplitudes as the ratios $M_{\lambda_1 \lambda_2}/M_{+0}^{even}$

$$\frac{M_{+0}}{M_{+0}^{even}} = 1 - \frac{24\pi^2 f_\gamma f_V^T m_V}{3f_V q^2} \left(\ln \frac{1-u_{min}}{u_{min}} - 2(1-2u_{min}) \right) \quad (4)$$

$$\frac{M_{++}}{M_{+0}^{even}} = \frac{m_V}{|q|\sqrt{2}} \left(\ln \frac{1-u_{min}}{u_{min}} - 1 + 2u_{min} \right) + \frac{24\pi^2 f_\gamma f_V^T}{3f_V |q|\sqrt{2}} \quad (5)$$

$$\frac{M_{+-}}{M_{+0}^{even}} = \frac{3m_V}{\sqrt{2}|q|} - \frac{48\pi^2 f_\gamma f_V^T m_V^2}{3f_V |q|^3 \sqrt{2}} \left(2 \ln \frac{1-u_{min}}{u_{min}} - 3(1-2u_{min}) \right) \quad (6)$$

The chiral-odd contributions to the amplitudes are accompanied with large numerical coefficients. In the case of M_{++} the chiral-even and the chiral-odd parts of (5) add with the same signs. In contrast, the chiral-even and the chiral-odd parts of M_{+0} or M_{+-} enters with opposite sign which leads to an effective

reduction of these amplitudes for intermediate $|t|$. Though the chiral-even and the chiral-odd contributions to M_{++} are of the same order with respect to $1/|\mathbf{q}|$ counting the chiral-odd one can be dominant up to very large $|t|$. According to (5) for the t range $3 \div 8 \text{ GeV}^2$ the chiral-even part constitutes only $10 \div 20\%$ of the M_{++} amplitude and for $|t| \approx 100 \text{ GeV}^2$ $M_{++}^{even} \approx 0.72 M_{++}^{odd}$. Due to a large compensation between chiral-even and chiral-odd parts of M_{+0} and M_{+-} the non spin-flip M_{++} amplitude dominates strongly in the region of intermediately high $|t|$. M_{+0} will exceed M_{++} only at $|t| > 40 (\text{GeV})^2$. For the $|t|$ interval $3 \text{ GeV}^2 \div 8 \text{ GeV}^2$

$$\frac{M_{+0}}{M_{++}} \sim 0.25 \div 0.35, \quad \frac{M_{+-}}{M_{++}} \sim -0.02 \div 0.04 \quad (7)$$

Both chiral-even and chiral-odd parts of the amplitudes were calculated in the leading order of $1/|\mathbf{q}|$ expansion. As usual in the QCD approach to any exclusive reaction, the question about the region of applicability of these results is open until the power corrections have not studied. In our case the situation is more difficult because the factorization of the amplitude into hard and soft parts is violated for the chiral-even part of M_{++} and the chiral-odd parts of M_{+0} and M_{+-} . In these cases the corresponding integrals over the quark longitudinal momentum u contain the end point logarithmic singularities. At present we simply restrict the corresponding u integrals to the interval $[1 - u_{min}, u_{min}]$, $u_{min} = m_V^2/\mathbf{q}^2$, which corresponds to the contribution of the hard region only. The good news is, however, that the chiral-odd part M_{++}^{odd} of the dominant in the intermediately large $|t|$ region helicity amplitude M_{++} is free from this end-point singularity. This chiral-odd part is numerically considerably larger than the hard contribution to its chiral-even counterpart M_{++}^{even} . Therefore we can expect that the relative uncertainty related with the uncalculated soft contributions to M_{++}^{even} is small.

Acknowledgments

This work was supported by the Foundation Universities of Russia (grant 015.02.01.16).

References

1. D.Yu. Ivanov, R. Kirschner, A. Schafer and L. Szymanowski, *Phys. Lett. B* **478**, 101 (2000).
2. B.L. Ioffe and A.V. Smilga, *Nucl. Phys. B* **232**, 109 (1984);
V.M. Belyaev and Y.I. Kogan, *Yad. Fiz.* **40**, 1035 (1984).